

Rational solutions of KZ equation, case S_4 .

Lev Sakhnovich

735 Crawford ave., Brooklyn, 11223, New York, USA.

E-mail address: lev.sakhnovich@verizon.net

Abstract

We consider the Knizhnik-Zamolodchikov system of linear differential equations. The coefficients of this system are generated by elements of the symmetric group S_n . We separately investigate the case S_4 . In this case we solve the corresponding KZ-equation in the explicit form.

Mathematics Subject Classification (2000). Primary 34M05, Secondary 34M55, 47B38.

Keywords. Symmetric group, natural representation, linear differential system, rational fundamental solution.

1 Introduction

1. We consider the Knizhnik-Zamolodchikov differential system (see [3])

$$\frac{dW}{dz} = \rho A(z)W, \quad z \in C, \quad (1.1)$$

where $A(z)$ and $W(z)$ are $n \times n$ matrices. We suppose that $A(z)$ has the form

$$A(z) = \sum_{k=1}^s \frac{P_k}{z - z_k}, \quad s \geq 2, \quad (1.2)$$

where $z_k \neq z_\ell$ if $k \neq \ell$. The matrices P_k are connected with matrix representation of the symmetric group S_n and are defined by formulas (2.1)-(2.4). A. Chervov and D. Talalaev formulated the following interesting conjecture [2].
Conjecture 1.1. *The Knizhnik-Zamolodchikov system (1.1), (1.2) has a rational fundamental matrix solution when parameter ρ is integer.*

We have proved this conjecture [5] for the case when $\rho = \pm 1$. In the present paper we separately consider the case S_4 and solve the corresponding KZ-equation in the explicit form.

2. In a neighborhood of z_k the matrix function $A(z)$ can be represented in the form

$$A(z) = \frac{a_{-1}}{z - z_k} + a_0 + a_1(z - z_k) + \dots \quad (1.3)$$

where a_k are $n \times n$ matrices. We investigate the case when z_k is either a regular point of $W(z)$ or a pole. Hence the following relation

$$W(z) = \sum_{p \geq m} b_p(z - z_k)^p, \quad b_m \neq 0 \quad (1.4)$$

is true. Here b_p are $n \times n$ matrices. We note that m can be negative.

Proposition 1.1. (necessary condition, see [4]) *If the solution of system (1.1) has form (1.4) then m is an eigenvalue of a_{-1} .*

We denote by M the greatest integer eigenvalue of the matrix ρa_{-1} . Using relations ((1.3) and (1.4)) we obtain the assertion.

Proposition 1.2. (necessary and sufficient condition, see [4]) *If the matrix system*

$$[(q+1)I_n - a_{-1}]b_{q+1} = \sum_{j+\ell=q} \rho a_j b_\ell, \quad (1.5)$$

where $m \leq q + 1 \leq M$ has a solution $b_m, b_{m+1}, \dots, b_M$ and $b_m \neq 0$ then system (1.1) has a solution of form (1.4).

We introduce the matrices

$$P_k^- = I + P_k, \quad P_k^+ = I - P_k. \quad (1.6)$$

2 Explicit solution

1. We consider the natural representation of the symmetric group S_n (see [1]). By $(i; j)$ we denote the permutation which transposes i and j and preserves all the rest. The $n \times n$ matrix which corresponds to $(i; j)$ is denoted by

$$P(i, j) = [p_{k,\ell}(i, j)], \quad (i \neq j). \quad (2.1)$$

The elements $p_{k,\ell}(i, j)$ are equal to zero except

$$p_{k,\ell}(i, j) = 1, \quad (k = i, \ell = j); \quad p_{k,\ell}(i, j) = 1, \quad (k = j, \ell = i), \quad (2.2)$$

$$p_{k,k}(i, j) = 1, \quad (k \neq i, k \neq j). \quad (2.3)$$

Now we introduce the matrices

$$P_k = P(1, k + 1), \quad 1 \leq k \leq n - 1. \quad (2.4)$$

and the $1 \times (n - 1)$ vector $e = [1, 1, \dots, 1]$, and $n \times n$ matrix

$$S = \begin{bmatrix} 2 - n & e \\ e^T & 0 \end{bmatrix}. \quad (2.5)$$

Using relations (1.38) and (2.1)-(2.4) we deduce that

$$T = (n - 2)I_n + S. \quad (2.6)$$

The eigenvalues of T are defined by the equalities

$$\lambda_1 = n - 1, \quad \lambda_2 = n - 2, \quad \lambda_3 = -1. \quad (2.7)$$

Hence we have $m_T = -1$, $M_T = n - 1$. We shall consider the case when $\rho = -1$. In this case the matrix solution $W(z)$ of system (1.1) can be written in the form

$$W(z) = \sum_{k=1}^s \frac{L_k}{z - z_k} + Q(z), \quad (2.8)$$

where L_k are $n \times n$ matrices, $Q(z)$ is a $n \times n$ matrix polynomial of the form

$$Q(z) = Q_{-1}z + Q_0. \quad (2.9)$$

By substituting formulas (2.8) and (2.9) in relation (1.1) we receive the equalities

$$(I - P_k)L_k = 0, \quad 1 \leq k \leq s, \quad (2.10)$$

$$\sum_{j \neq k} \frac{P_k L_j + P_j L_k}{z_k - z_j} + P_k Q(z_k) = 0, \quad (2.11)$$

$$Q_{-1} = -TQ_{-1}. \quad (2.12)$$

2. Further we consider the case S_4 .

The vectors L_k can be chosen in the following form

$$L_1 = \text{col}[1, 1, -1, -1], \quad L_2 = \alpha \text{col}[1, -1, 1, -1], \quad L_3 = \beta \text{col}[1, -1, -1, 1]. \quad (2.13)$$

We note, that L_k which are defined by relation (2.13) satisfy conditions (2.10). The numbers α and β are defined by relations

$$\alpha = -\frac{z_3 - z_2}{z_3 - z_1}, \quad \beta = \frac{z_3 - z_2}{z_2 - z_1}. \quad (2.14)$$

Now we introduce the matrix polynomial $Q(z)$ (see (2.9)), where

$$Q_{-1} = -[(z_2 - z_1)(z_3 - z_1)]^{-1} \text{col}[3, -1, -1, -1], \quad (2.15)$$

$$Q_0 = [(z_2 - z_1)(z_3 - z_1)]^{-1} (z_1 N_1 + z_2 N_2 + z_3 N_3). \quad (2.16)$$

Here the vectors N_k are defined by relations

$$N_1 = L_1 \quad N_2 = \text{col}[1, -1, 1, -1], \quad N_3 = \text{col}[1, -1, -1, 1]. \quad (2.17)$$

By direct calculations we see that that matrices L_k , ($k = 1, 2, 3$) and matrix polynomial $Q(z)$ which are defined by relations (2.9) and (2.13)-(2.17), satisfy the relations (2.10)-(2.12). Hence the vector function

$$Y_1(z) = \sum_{k=1}^3 \frac{L_k}{z - z_k} + Q(z) \quad (2.18)$$

is the rational solution of system (1.1) when $\rho = -1$.

In order to construct the second rational solution of system (1.1) we introduce the vector

$$M_k = \beta_k \text{col}[1, 1, 1, 1], \quad k = 1, 2, 3, \quad (2.19)$$

where the numbers β_k are defined by the relations

$$\beta_1 = z_2 - z_3, \quad \beta_2 = z_3 - z_1, \quad \beta_3 = z_1 - z_2. \quad (2.20)$$

By direct calculations we see that the numbers β_k satisfy the equations

$$\frac{\beta_1 + \beta_2}{z_1 - z_2} + \frac{\beta_1 + \beta_3}{z_1 - z_3} = 0, \quad (2.21)$$

$$\frac{\beta_1 + \beta_2}{z_2 - z_1} + \frac{\beta_2 + \beta_3}{z_2 - z_3} = 0. \quad (2.22)$$

It follows from relations (2.19)-(2.22) that conditions (2.10)-(2.12) will be fulfilled if $L_k = M_k$, $Q_{-1} = Q_0 = 0$. Hence the vector function

$$Y_2(z) = \sum_{k=1}^3 \frac{M_k}{z - z_k} \quad (2.23)$$

is the rational solution of system (1.1) when $\rho = -1$.

In order to construct the next solution $Y_3(z)$ of system (1.1) we introduce the vectors

$$m_1 = \beta_1 \text{col}[0, 0, 1, a], \quad m_2 = \beta_2 \text{col}[0, b, 0, c], \quad m_3 = \beta_3 \text{col}[0, 1, a, 0], \quad (2.24)$$

where

$$a = -\frac{\beta_1}{\beta_3}, \quad b = -\frac{\beta_3}{\beta_2}, \quad c = \frac{\beta_1^2}{\beta_2\beta_3}. \quad (2.25)$$

We note, that the numbers β_k as in the previous case are defined by relations (2.20). It follows from relations (2.24) and (2.25) that conditions (2.10)-(2.12) will be fulfilled if $L_k = m_k$, $Q_{-1} = Q_0 = 0$. Hence the vector function

$$Y_3(z) = \sum_{k=1}^3 \frac{m_k}{z - z_k} \quad (2.26)$$

is the rational solution of system (1.1) when $\rho = -1$.

In order to construct the next solution $Y_4(z)$ of system (1.1) we introduce the vectors

$$\ell_1 = \alpha_1 \text{col}[0, 0, 1, a], \quad \ell_2 = \alpha_2 \text{col}[0, b, 0, c], \quad \ell_3 = \alpha_3 \text{col}[0, d, e, 0], \quad (2.27)$$

where

$$a = -\frac{\alpha_1}{\alpha_3}, \quad b = -\frac{\alpha_3}{\alpha_2}d, \quad c = \frac{\alpha_1^2}{\alpha_2\alpha_3}, \quad (2.28)$$

$$d = -\frac{\alpha_1^2}{\alpha_2\alpha_3^3}(\alpha_1\alpha_2 + \alpha_3^2), \quad e = 1 + \frac{\alpha_1}{\alpha_2}. \quad (2.29)$$

We note, that the numbers α_k are defined by the relations

$$\alpha_1 = (z_3 - z_2)^{-1}, \quad \alpha_2 = (z_1 - z_3)^{-1}, \quad \alpha_3 = (z_2 - z_1)^{-1}. \quad (2.30)$$

It follows from relations (2.27)-(2.30) that conditions (2.10)-(2.12) will be fulfilled if $L_k = \ell_k$, $Q_{-1} = Q_0 = 0$. Hence the vector function

$$Y_4(z) = \sum_{k=1}^3 \frac{\ell_k}{z - z_k} \quad (2.31)$$

is the rational solution of system (1.1) when $\rho = -1$.

Proposition 2.1. *Let us consider the case S_4 when $\rho = -1$. In this case the fundamental solution of system (1.1) has the form*

$$Y(z) = \sum_{k=1}^4 c_k Y_k(z), \quad (2.32)$$

where c_k are arbitrary constants, the functions $Y_k(z)$ are defined by relations (2.18), (2.23), (2.26) and (2.31).

Remark 2.1. The explicit solution for the case S_3 was constructed in the paper [5].

References

1. Burrow M., Representation Theory of Finite Groups, Academic Press, 1965.
2. Chervov A., Talalaev D., Quantum Spectral Curves, Quantum Integrable Systems and the Geometric Langlands Correspondence, arXiv:hep-th/0604128, 2006.
3. Etingof P.I., Frenkel I.B., Kirillov A.A. (jr.), Lectures on Representation Theory and Knizhnik-Zamolodchikov Equations, Amer. Math. Society, 1998.

4. Sakhnovich L.A., Meromorphic Solutions of Linear Differential Systems, Painleve Type Functions, arXiv:math CA/0607555,2006
5. Sakhnovich L.A., Rational Solutions of KZ equation, existence and construction, arXiv.math:-ph/0609067, 2006.